

Moxifloxacin Eye Drops IP



लूमीमोक्स आई ड्रॉप्स

Composition

Each ml. Contains :
Moxifloxacin Hydrochloride IP
Equivalent to Moxifloxacin 5.0 mg
Sterile aqueous vehicle q.s.
Preservative : Product is self preserved

Category:

Moxifloxacin as antibiotic.

Pharmacology:

Mode of action

LumeMOX constitutes 4th generation Fluoro-Quinolones, Moxifloxacin as an agent. Floro-Quinolones exert their antibacterial effect by preventing bacterial DNA from unwinding and duplicating. Moxifloxacin hydrochloride is a synthetic 8-methoxyfluoroquinolone with a broad spectrum anti-bacterial effect on gram positive, gram negative and other anaerobic micro-organism.

It is "PRESERVATIVE FREE" providing freedom from

hypersensitivity or allergy associated with the preservatives. Well tolerated with minimal ocular side effects provides increased penetration into ocular tissues and fluids with improved activity against Streptococci and Staphylococci species. It has a moderate to excellent activity against clinically relevant, gram-negative ocular pathogens.

Indications and Usage:

Bacterial Conjunctivitis (pink eye), Bacterial Corneal Ulcer Infection, caused by the following susceptible strains-

Micrococcus luteus	Staphylococcus warneri
Staphylococcus aureus	Streptococcus pneumoniae
Staphylococcus epidermidis	Streptococcus viridans group
Staphylococcus haemolyticus	Acinetobacter Iwoffii
Staphylococcus hominis	Haemophilus influenzae
Haemophilus parainfluenzae	Chlamydia trachomatis

Precautions:

Hypersensitivity towards any of the component in the solution.

Warning:

- Growth of Resistant Organisms with Prolonged Use

- Avoidance of Contact Lens Wear
- if irritations persist, discontinue the use and consult the eye doctor.

Adverse effects:

Nausea, Diarrhea, Dizziness, blurred vision that does not go away, burning, stinging, or itching of the eyes or eyelids. Ocular hyperemia, Ocular pain and Ocular pruritus

Dosage and Administration:

A drop in the affected eyes twice a day or as directed by your doctor.

Information to Patients:

Shake well before use.

Follow the instructions that are provided by your doctor and if the adverse effect persists, discontinue the usage and contact your doctor immediately.

This drug may cause low blood sugar and mental health related side effects.

Mfg. Lic. No.: 78/UA/SC/P-2006

Marketed by: Lumera
(A Division of Axa Parenterals Ltd.)

Manufactured in India by:
Axa Parenterals Ltd.
Plot No.: 936, 937 & 939
Kishanpur Jamalpur
Roorkee-247 667, Uttarakhand